

POSTER PRESENTATION

Open Access

# Inhibitory effect of novel iron chelator, 1-(N-acetyl-6-aminohexyl)-3-hydroxy-2-methylpyridin-4-one (CM1) and green tea extract on growth of *Plasmodium falciparum*

Somdet Srichairatanakool<sup>1\*</sup>, Supalerk Thipubol<sup>1</sup>, Wachiraporn Tipsuwan<sup>2</sup>, Chairat Uthaipibull<sup>3</sup>

From Challenges in malaria research: Core science and innovation  
Oxford, UK. 22-24 September 2014

## Background

Iron is an essential micronutrient required by all living organisms including malaria parasites (*Plasmodium* spp.) for many biochemical reactions, especially growth and multiplication processes. Therefore, the malaria parasites needs to take up the iron from outside or/and inside the parasitized red blood cells (PRBC). Iron chelators are widely used for treatment of thalassemia-related iron overload and also inhibit parasite growth at levels which are non-toxic to mammalian cells.

## Materials and methods

The inhibitory effect of 1-(N-acetyl-6-amino-hexyl)-3-hydroxypyridin-4-one (CM1) and green tea extract (GTE) on the growth of malaria parasite *P. falciparum* was investigated compared with standard chelators including desferrioxamine (DFO), deferiprone (DFP) and deferasirox (DFX). A flow cytometric technique was used to enumerate the PRBC stained with SYBR Green I fluorescent dye. Labile iron pool (LIP) was assayed using calcein-acetomethoxy (Calcein-AM) fluorescent method.

## Results

The IC<sub>50</sub> values of DFO, GTE, CM1, DFX and DFP against *P. falciparum* were 14.09, 21.11, 35.14, 44.71 and 58.25 μM, respectively. Importantly, CM1 was more effective in reducing LIP level in the *P. falciparum* culture than DFP ( $P < 0.05$ ).

## Conclusions

CM1 and GTE exhibit anti-malarial activity. They could interfere with uptake of exogenous iron or deplete intracellular labile iron in malaria parasites, leading to inhibition of their growth.

## Acknowledgements

The work has been supported by the Thailand Graduate Institute of Science and Technology (TGIST), National Science and Technology Development Agency (NSTDA) and Faculty of Medicine, Chiang Mai University.

## Authors' details

<sup>1</sup>Chiang Mai University, Chiang Mai, Thailand. <sup>2</sup>University of Phayao, Phayao, Thailand. <sup>3</sup>National Science and Technology Development Agency, Pathum Thani, Thailand.

Published: 22 September 2014

doi:10.1186/1475-2875-13-S1-P84

**Cite this article as:** Srichairatanakool et al.: Inhibitory effect of novel iron chelator, 1-(N-acetyl-6-aminohexyl)-3-hydroxy-2-methylpyridin-4-one (CM1) and green tea extract on growth of *Plasmodium falciparum*. *Malaria Journal* 2014 **13**(Suppl 1):P84.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Chiang Mai University, Chiang Mai, Thailand  
Full list of author information is available at the end of the article